

FIGURE 4 | bFGF + NSCs significantly improved behavioral recovery after stroke.







FIGURE 2




FIGURE 3

## Human bFGF amino acid sequences

### **22.5 kD form (Genbank GI: 482272; SEQ. ID. No. 1)**

1 mdrgrgral pgrlggrgr grapervggr grgrgtaapr aapaargsrp gpagtmaags  
61 ittlpalped ggsgafppgh fkdprlyck nggfflrhp dgrvdgvrek sdphiklqlq  
121 aeergvvsik gvcanrylam kedgrllask cvtdecfffe rlesnnynty rsrkytswyv  
181 alkrtgqykl gsktgpgqka ilflpmsaks

### **114 amino acid form (Genbank GI: 8250666; SEQ. ID. No. 2)**

1 ldrgrgral pgrlggrgr grapervggr grgrgtaapr aapaargsrp gpagtmaags  
61 ittlpalped ggsgafppgh fkdprlyck nggfflrhp dgrvdgvrek sdph

### **88 amino acid form (Genbank GI: 4261553; SEQ. ID. No. 3)**

1 flrhpdgrv dgvreksdph iklqlqaeer gvvsikgvca nrylamkedg rllaskcvtd  
61 ecfferles nnyntyrsk ytswyval